BRIEF REPORT

IDENTIFICATION BY PCR-FRLP OF BACTERIA PRESENT IN THE COLOSTRUM OF WOMEN Residing in Barranquilla, Colombia

Luz A. Sarmiento-Rubiano^{1,a}, Leidys Goenaga^{2,b}, Marianella Suarez-Marenco^{2,c}, Clara Gutierrez-Castañeda^{2,c}, Carmen M. Sarmiento^{3,d}, Jimmy Becerra Enríquez^{1,e}

¹ Food and Human Behavior Research Group, Metropolitan University, Barranquilla, Colombia.

² Ecological and Agro-industrial Management Research Group, Microbiology Program, Free University, Barranquilla, Colombia.

³ Preventive and Community Medicine, University of Sinú, Cartagena, Colombia.

^a Bacteriologist, PhD in Food Technology,^b bacteriologist, master's degree in Microbiology; ^cmicrobiologist, master's degree in Microbiology; ^d bacteriologist, PhD in Public Health; ^c biologist, PhD in Biomedicine and Biotechnology.

ABSTRACT

With the aim of isolating and identifying bacteria present in the colostrum of women in the city of Barranquilla, located in the Colombian Caribbean, we carried out a descriptive study on 55 colostrum samples, which were cultured on M17, MRS, and TOS agar and incubated under aerobic and anaerobic conditions. A total of 350 microorganisms were isolated, of which 296 were identified at the genus level by PCR-RFLP with the enzymes HaeIII and RsaI. The online program kodebio.shinyapps.io/RFLP-inator was used for the in-silico identification of the isolates. Seven hypothetical bacterial genera were identified: *Staphylococcus, Lactobacillus, Enterococcus, Bifidobacterium, Corynobacterium, Streptococcus,* and *Leuconostoc,* with *Staphylococcus, Lactobacillus, and Enterococcus* being the most dominant, representing 61.1% of the identified microorganisms. Our results are a starting point for understanding the microbial composition of human colostrum, considering the particular context and environmental conditions of the Colombian Caribbean.

Keywords: Colostrum; Polymorphism, Restriction Fragment Length; Microbiota, Breast Feeding (source: MeSH NLM).



Cite as: Sarmiento-Rubiano LA, Goenaga L, Suarez-Marenco M, Gutierrez-Castañeda C, Sarmiento CM, Becerra Enríquez J. Identification by PCR-FRLP of bacteria present in the colostrum of women residing in Barranquilla, Colombia. Rev Peru Med Exp Salud Publica. 2025;42(2):184-9. doi:10.17843/rpmesp.2025.422.14321.

Correspondence. Luz A. Sarmiento-Rubiano; lusarru@hotmail.com

Received. 11/09/2024 **Approved.** 22/05/2025 **Online.** 11/06/2025



This work is licensed under a Creative Commons Attribution 4.0 International

INTRODUCTION

Currently, more than 590 bacterial genera have been identified in human milk, belonging mainly to the *phyla Firmicutes, Actinobacteria*, and *Protebacteria*, with the genera *Staphylococcus, Streptococcus, Lactobacillus, Pseudomonas, Bifidobacterium, Corynebacterium, Enterococcus, Acinetobacter*, and *Clostridium*, among others, being particularly dominant ⁽¹⁾.

Human milk has all the components and nutrients necessary for the proper growth, development, and protection of newborns, and is their main source of commensal, mutualistic, and potentially probiotic bacteria ⁽²⁾. The bacteria provided by breast milk perform various functions, including: reducing the incidence and severity of infections ⁽³⁾; maturing the immune system ⁽⁴⁾; defending against pathogenic bacteria through the presence of bacterial phages ⁽⁵⁾; and exposure to the maternal milk microbiome promoting the intestinal health of the newborn and their physical and mental development ⁽⁶⁾.

The composition of the human milk microbiota varies according to the stage of lactation and factors such as the mother's hygiene and diet, her gut microbiota, antibiotic use, geographical location, lifestyle, and genetic, epigenetic, and ethnic factors ⁽⁷⁾. Colostrum is produced by the mammary glands from the moment of birth and for six days thereafter. It is rich in cytokines, antimicrobial peptides, antibodies, hormones, and various bioactive compounds ⁽²⁾. The colostrum microbiota is strongly associated with the maternal environment, as well as the type of delivery (natural or cesarean), and the evolution from colostrum to mature milk involves signi-

ficant changes in its microbiota, in which the interaction of the infant's emerging oral microbiome with the mammary gland is fundamental ⁽⁸⁾.

Studying microorganisms in human milk under different environmental and population conditions contributes to our knowledge of this fluid and its potential impact on the health of infants and mothers. Culture-dependent and culture-independent techniques allow the identification of a wide variety of bacteria, each with advantages and disadvantages that are evident in the obtained results. However, culture techniques allow the isolation and preservation of microorganisms with potential probiotic characteristics. This study aimed to isolate and identify bacteria from the colostrum of women in the city of Barranquilla, in the Colombian Caribbean, using the culture technique and subsequent identification by PCR-FRLP (Restriction Fragment Length Polymorphism).

THE STUDY

A descriptive cross-sectional study was conducted, including 55 postpartum women who were treated at the maternity ward of the Metropolitan Hospital in the city of Barranquilla (Colombian Caribbean) and who met the following inclusion criteria: being of legal age, having had a minimum of 37 weeks of gestation, vaginal delivery without complications or antibiotic use, being in good health, and having voluntarily agreed to participate by donating 3 to 5 ml of colostrum, collected between 12 and 48 hours postpartum. The colostrum was obtained after washing the nipples with mild soap and sterile water, using an electric breast pump (Medela Swing) under sterile conditions. The first two drops of colostrum were discarded, and 3 to 5 ml of sample were collected in a sterile container and stored at -80°C until analysis.

To isolate bacteria from colostrum, 100 µl of sample was directly seeded onto MRS (Man Rogosa Sharpe, for the isolation of acid-lactic bacteria), M17 (for lactic *Streptococcus*) and Propionate TOS agar (for bifidobacteria) culture media. The agar plates were incubated at 37 °C for 72 hours (MRS and M17 under aerobic and anaerobic conditions—Oxoid Anaerogen system and TOS under anaerobic conditions only). The colonies isolated from the different cultures were morphologically identified by Gram staining, re-cultured in liquid culture medium according to the agar of origin or in Brain Heart Infusion (BHI) broth, and cryopreserved at -80 °C with 20% glycerol for future studies.

The gender identification of bacteria isolated from colostrum was performed using PCR-RFLP. Approximately

KEY MESSAGES

Motivation for the study. There is little information on the microbiota found in the colostrum of women living in the Colombian Caribbean. Identifying and isolating microorganisms in this fluid is of interest to both medicine and industry.

Main findings. The dominant bacterial genera found in colostrum were *Staphylococcus, Lactobacillus,* and *Enterococcus.* Microorganisms with probiotic potential were isolated and should be studied in greater detail.

Public health implications. Understanding the microbiota found in colostrum allows us to understand its contribution to infant health and develop strategies to enhance the beneficial impact of these microorganisms on their development.

1508 base pairs of r16S DNA were amplified by colony PCR with universal primers 27F: 5'-AGAGTTTGATCCTGGCT-CAG-3' and 1492R: 5'-GGTTACCTTGTTACGACTT-3' ⁽⁹⁾. Each 50 μl PCR reaction contained: 1X buffer (BIOTOOLS), 1.5 mM MgCl2, 200 μM dNTPs, 20 pmol of each primer, 1U Taq polymerase (BIOTOOLS), and a small portion of colony added with a sterile toothpick. *Lactococcus lactis* NZ900 was used as a positive control for the reaction. The amplification program was as follows: five minutes at 96 °C; 35 cycles: 95 °C for 30 seconds, 60 °C for 30 seconds, 72 °C for one minute; and one cycle of 72 °C for seven minutes (BIO-RAD Thermocycler). The PCR products were visualized on 0.8% agarose gels stained with Syber-Green. Electrophoresis was performed with 1X TAE buffer at 80V for 30 minutes (Bio-Rad horizontal electrophoresis).

For RFLP, PCR amplificates were digested in separate reactions with the enzymes HaeIII (GC^GC) and RsaI (GT^AC) according to the manufacturer's recommendations (New England Biolab) (10 μ l of PCR product, 2 units of enzyme, 2 μ l 10X buffer in a final volume of 20 μ l). The reactions were incubated for 3 hours at 37 °C. The digestion products were subjected to electrophoresis in 2% agarose gels with Syber-Green and the generated band profile was visualized on a Biorrad ChemiDocTM XRS photodocumenter. The band sizes obtained were compared with the HyperLadderTM 100pb DNA–Bioline molecular weight marker.

The resulting band profiles were classified into eight groups of microorganisms that showed identical restriction profiles with the two enzymes. Using the online program kodebio.shinyapps.io/RFLP-inator, which allows bacterial identification based on *in silico* simulation of 16S DNA restriction profiles, it was possible to identify the isolates obtained at the genus level. *Lactococcus lactis* subsp. cremoris NZ9000 (GenBank NCBI sequence NC_017949.1) was used as a positive control, whose restriction profiles with HaeIII and RsaI were verified *in silico*.

We conducted a descriptive statistical analysis. Microbial growth is expressed as log¹⁰ CFU/ml. The identification of microorganisms by PCR-RFLP is reported as relative frequencies.

The research protocol for this study was approved by the Ethics Committee of the Metropolitan University (code 0340517) and all participants signed an informed consent form in accordance with Resolution 8430 of 1993 of the Colombian Ministry of Health and Social Protection and the Declaration of Helsinki.

FINDINGS

Of the 55 colostrum samples cultured, only 90.9% (n=50) showed microbial growth in any of the culture media under the described conditions. The average number of microorganisms isolated from each colostrum was 2.73 log¹⁰ CFU/ml (range 1.30 to 3.47). Microbial growth in each of the culture media and evaluated conditions had an average value of 2.39 log¹⁰ CFU/ml (Table 1). Taking into account the different culture conditions and differential morphology of the co-lonies, it was possible to isolate 350 microorganisms. Gram staining was used to determine the microscopic morphology of the isolates; in M17 medium, mainly Gram-positive cocci were observed, in MRS medium, Gram-positive coccobacilli and Gram-positive bacilli, and in TOS medium, Gram-positive cocci were cocci and coccobacilli.

PCR amplification (\approx 1508 bp) of r16S DNA was possible in all the isolated microorganisms. Digestion with the enzymes *HaeIII* and *RsaI* resulted in 332 and 335 band profiles. Since not all isolates showed a defined pattern (diffuse bands, imperceptible bands, or non-digestion of the DNA fragment), only 296 isolates were selected for identification (Figure 1). Since many of the bacteria presented the same band profile with both enzymes, the profiles were grouped, obtaining eight restriction patterns from the 296 microorganisms (Table 2).

Bacterial genera were identified based on their growth in the culture medium, Gram staining morphology, and the results of *in silico* band pattern analysis using the program kodebio.shinyapps.io/RFLP-inator.

The highest number of microorganisms (57.2%, n=200) was isolated and identified from the M17 culture medium. We identified bacteria of the genus *Staphylococcus* (patterns 1 and 2) growing under aerobic and anaerobic conditions, which showed a band profile common to digestion with *HaeIII*, but different with *RsaI*; and bacteria of the genus *Enterococcus* (pattern 3). Microorganisms grown on M17 agar only under anaerobic conditions accounted for 8.2% of the isolates, which were identified as bacteria of the genus *Streptococcus* (pattern 4).

We found that 22.8% of the identified microorganisms were isolated in the MRS medium, under aerobic and anaerobic conditions. *Lactobacillus* (pattern 5) was identified, as well as *Leuconostoc* (pattern 6), which grew only under anaerobic conditions. The bacteria recovered and identified in the TOS medium under anaerobic conditions (band patterns 7 and 8), representing 20% of the isolates, belonged to the genera *Corynebacterium* sp. and *Bifidobacterium*, respectively (Table 2).

DISCUSSION

In this study, seven hypothetical bacterial genera were isolated and identified by PCR-RFLP from 50 human colostrum samples, belonging to the genera *Staphylococcus*, *Lactobacillus*, *Enterococcus*, *Bifidobacterium*, *Corynobacterium*,

Table 1. Number of microorganisms isolated from colostrum samples from each culture medium and different conditions.

Culture medium	Condition	Samples with growth	Average	Number of isolated	
		n=50 (%)	log ¹⁰ CFU /ml	microorganisms (n=550)	
1417	Aerobic	40 (80)	2.42	128	
M11/	Anaerobic	28 (56)	2.27	72	
MDC	Aerobic	16 (32)	2.41	53	
MRS	Anaerobic	8 (16)	2.59	27	
TOS	Anaerobic	23 (46)	2.36	70	

CFU: colony-forming units, M17: M17 agar culture medium, MRS: Man Rogosa Sharpe medium, TOS: propionate agar medium.



^a Shows the causes of exclusion of microorganisms in the process.

Figure 1. Flowchart of the process of culturing and identifying microorganisms in human colostrum

Streptococcus, and *Leuconostoc*, with *Staphylococcus*, *Lactobacillus* and *Enterococcus*, corresponding to 61.1% of the 296 microorganisms identified from 350 isolates.

Other authors, who used culture-dependent methods, have identified Staphylococcus, Streptococcus, Corynobacterium, and Cutibacterium as the dominant bacterial genera in human milk, and in smaller proportions Lactococcus, Enterococcus, Lactobacillus, Leuconostoc, Weisella, and Bifidobacterium^(2,10). As in this study, Staphylococcus-predominant milk flora has been previously reported. For example, in women from urban areas of China, r16S DNA sequencing showed that more than 40% of the milk microbiota corresponded to bacteria of the genera Staphylococcus and Streptococcus⁽¹¹⁾. A meta-analysis that included twelve studies from different parts of the world found that the dominant genera in milk, regardless of geographical location, were Staphylococcus and Streptococcus (12). In this regard, it should be noted that the microbiota evolves from colostrum to mature milk and that studies such as that by Jiménez et al. in 2008, in colostrum from 36 Spanish women, found Staphylococcus and Enterococcus to be the dominant genera, with a lower proportion of Lactobacillus (13). These results are consistent with those found in this study, in which Enterococcus and Lactobacillus were the most frequent genera after Staphylococcus.

The mother's diet, diabetes, obesity, and even the sex of the baby, among other factors, have been identified as influencing the composition of the colostrum microbiota ⁽¹⁴⁾. Xie *et al.*, evaluating 97 healthy mothers on the Chinese island of Hainan, found that the ethnic origin of individuals may be an important factor associated with the diversity of the colos-

trum microbiome $^{\scriptscriptstyle (15)}$. In Colombia, and specifically in the Caribbean, there is great ethnic and cultural diversity characterized by a mixture of African, European, and Native American ancestry (16). This diversity gives rise to such varied ethnic and genetic profiles that it would be very difficult to establish relationships between the milk microbiota and ethnicity, especially when only gender identification of the species found in colostrum is available. With regard to maternal diet, it is known that the consumption of macronutrients, mainly fat, modifies the nutritional quality of colostrum ⁽¹⁷⁾. The availability of nutrients in both the milk fluid and the infant's intestinal mucosa affects the variety and quantity of existing bacterial populations. In this study, maternal diet was not evaluated. In Colombia, there are few studies on human milk microbiota. Among them, Londoño-Sierra et al., evaluating 101 lactating women in the city of Medellín, demonstrated a direct relationship between nutritional status and maternal diet with the dominant microbial genera in milk (18).

The number of bacteria present in human milk reported by other authors range between 1.5 and 4.0 log¹⁰ CFU/ ml, with higher values in colostrum, between 3.14–4.22 log¹⁰ CFU/ml, which can increase to 6 log¹⁰ CFU/mL in cases of mastitis ⁽¹⁹⁾. In this study, we found a lower number of microorganisms in the samples, 2.73 log¹⁰ CFU/ml (range 1.30 to 3.47), a reduction associated with the fact that only microorganisms cultivable under the conditions described were included. There is no doubt that the application of molecular techniques based on the massive sequencing of r16S

			Identification of band profiles by RFLP-inator			% MI
Pattern	Culture media	Gram stain	*Hae III	[∗] Rsa I	Hypothetical microorganisms	(n=296)
1	M17 aerobiosis M17 anaerobiosis	Gram + Cocci	1200, 310	630, 490, 380	HaeIII: S004065075 Staphylococcus aureus 08BA02176 RsaI: S000404448 uncultured rumen bacterium BE38	16.4
2	M17 aerobiosis M17 anaerobiosis	Gram + Cocci	1200, 310	500, 490, 410, 120	HaeIII: S004443882 Staphylococcus epidermidis SEI RsaI: S001095749 Staphylococcus epider- midis SSL08	20.2
3	M17 anaerobiosis M17 anaerobiosis	Gram + Cocci	600, 460, 290, 120, 30	900, 350, 150, 120	HaeIII: S000892225 Enterococcus mund- tii HDYM-22 RsaI: S002949963 Enterococcus sp. PPC34A	12.4
4	M17 anaerobiosis	Gram + Cocci	580 450, 310, 120	620, 360, 260, 150, 120	HaeIII: S000470127 Streptococcus sali- varius subsp. null DL1 RsaI: S002032101 Streptococcus sp.	8.2
5	MRS aerobiosis MRS anaerobiosis	Gram + bacilli	570, 460, 330, 120, 30	900, 350, 140, 110	HaeIII: S003260210 Lactobacillus pen- tosus ZU 28 RsaI: S003260206 Lactobacillus pentosus ZU 24	15.1
6	MRS anaerobiosis	Gram + Cocci- bacilli	1100, 270, 70, 45	490, 400, 120, 100	HaeIII: S000430949 Leuconostoc pseu- domesenteroides RO1 RsaI: S000978801 Leuconostoc pseudo- mesenteroides QL28	7.7
7	TOS anaerobiosis	Gram + bacilli	460, 360, 160, 95, 85, 65, 55, 35	410-210-160- 150-115-85	HaeIII: S001576581 Corynebacterium sp. ICIRC105 RsaI: S000721364 Corynebacterium aurimucosum AE1-3	8.5
8	TOS anaerobiosis	Gram + bacilli	260, 200, 150, 100, 90, 70, 50, 30, 20	630, 360, 250, 150, 120	HaeIII: S000871434 Bifidobacterium animalis subsp. lactis IDCC 4301 RsaI: S000871434 Bifidobacterium animalis subsp. lactis IDCC 4301	11.5
Control	M17 anaerobiosis	Gram + Diplococci	457, 410, 309, 187, 123, 22	890, 355, 146, 83, 34	<i>Lactococcus lactis</i> subsp. cremoris NZ9000, (NC_017949.1)	

Table 2. Identification of bacterial genera isolated from colostrum samples using PCR-RFLP.

The band profiles obtained with each digestion enzyme are shown according to the band sizes generated.

^a HaeIII and RsaI are restriction enzymes used in digestion.

MI: identified microorganisms. RFLP: Restriction Fragment Length Polymorphism, M17: Agar M17 culture medium, MRS: Man Rogosa Sharpe medium, TOS: propionate agar medium.

ribosomal subunit genes allows for the accurate assessment of the microbial diversity of human milk. However, culture-based techniques may be a viable alternative for studying the milk microbiota when sequencing techniques are not available. Tabit *et al.*, in their review of molecular techniques in the study of milk, report that PCR-RFLP is one of the methodologies with the highest reproducibility of results, although it does not have much discriminatory power ^{(20).}

The main limitations of this study should be considered in order to interpret our results, these being mainly the small sample size, the lack of randomization in its selection, and the absence of information regarding the mother's diet as a determining factor of the microbiota. The use of culture techniques in the study of the microbiota limits the findings to cultivable microorganisms. The usefulness of PCR-RFLP for identifying microbial isolates only allows for identification at the genus level.

In conclusion, the findings of our study constitute a starting point in understanding the microbial composition of human colostrum, considering the particular context and environmental conditions of the Colombian Caribbean. These results are consistent with reports from other parts of the world, which point to the dominance of genera such as *Staphylococcus, Lactobacillus,* and *Enterococcus*. It is important to implement mass sequencing techniques to consolidate this finding and achieve species- and subspecies-level identification of the microorganisms present in colostrum. The identification and characterization of potential probiotics among the isolated microorganisms is of great interest to medicine and industry.

Acknowledgments. The authors would like to thank the breastfeeding mothers for donating their milk samples, as well as the Fundación Hospital Metropolitano de Barranquilla for facilitating access to their patients.

Author contributions. All authors declare that they comply with the authorship criteria recommended by the ICMJE.

CRediT roles. LAS: research, visualization, writing—original draft, software, and validation. JEB: project management, conceptualization,

REFERENCES

- Zimmermann P, Curtis N. Breast milk microbiota: A review of the factors that influence composition. J Infect. 2020;81(1):17-47. doi: 10.1016/j.jinf.2020.01.023.
- El-Loly M. Colostrum ingredients, its nutritional and health benefits-an overview. Clin Nutr Open Sci. 2022;44:126-143. doi: 10.1016/j.nutos.2022.07.001.
- Fernández L, Pannaraj PS, Rautava S, Rodríguez JM. The microbiota of the human mammary ecosystem. Front Cell Infect Microbiol. 2020;10:586667. doi: 10.3389/fcimb.2020.586667.
- Lyons KE, Ryan CA, Dempsey EM, Ross RP, Stanton C. Breast milk, a source of beneficial microbes and associated benefits for infant health. Nutrients. 2020;12(4):1039. doi: 10.3390/nu12041039.
- Lugli GA, Milani C, Turroni F, Tremblay D, Ferrario C, Mancabelli L, et al. Prophages of the genus Bifidobacterium as modulating agents of the infant gut microbiota. Environ Microbiol. 2016;18:2196–2213. doi: 10.1111/1462-2920.13154.
- Granger CL, Embleton ND, Palmer JM, Lamb CA, Berrington JE, Stewart CJ. Maternal breastmilk, infant gut microbiome and the impact on preterm infant health. Acta Paediatri. 2021;110(2):450-457. doi: 10.1111/apa.15534.
- Moossavi S, Sepehri S, Robertson B, Bode L, Goruk S, Field CJ, *et al.* Composition and variation of the human milk microbiota are influenced by maternal and early-life factors. Cell Host Microbe. 2019;25:324–335.e4. doi: 10.1016/j.chom.2019.01.011.
- Toscano M, De Grandi R, Peroni DG, Grossi E, Facchin V, Comberiati P, et al. Impact of delivery mode on the colostrum microbiota composition. BMC Microbiology. 2017;17:205. doi: 10.1186/s12866-017-1109-0.
- Lee H, Lee CK, Kim K. Isolation of Novel Strains of Lactobacillus gasseri EJL and Bifidobacterium breve JTL from Breast Milk and Infant Feces: A Longitudinal Study of a Mother-infant Pair. Microbiol Biotechnol Lett. 2021; 49(1): 1-8. doi: 10.48022/mbl.2010.10011.
- Hunt KM, Foster JA, Forney LJ, Schütte UME, Beck DL, Abdo Z, et. al. Characterization of the diversity and temporal stability of bacterial communities in human milk. PLoS One. 2011;6:1–8. doi: 10.1371/journal.pone.0021313.
- Sakwinska O, Moine D, Delley M, Combremont S, Rezzonico E, Descombes P, *et al.* Microbiota in breast milk of Chinese lactating mothers. PLoS One. 2016;11(8):e0160856. doi: 10.1371/journal.pone.0160856.

and methodology. LG, MSM: research, data curation, visualization, and methodology. CGC: formal analysis, research, software, and fund acquisition. MCS: writing—review and editing.

Funding. This work has been supported by the Metropolitan University. All experimental work was carried out in the institution's laboratories and with its resources.

Conflicts of interest. The authors declare that they have no conflicts of interest.

- Fitzstevens JL, Smith KC, Hagadorn JI, Caimano MJ, Matson AP, Brownell EA. Systematic review of the human milk microbiota. Nutr Clin Pract. 2017;32(3):354-364. doi: 10.1177/0884533616670150.
- Jiménez E, Delgado S, Fernández L, García N, Albújar M, Gómez A, et al. Assessment of the bacterial diversity of human colostrum and screening of staphylococcal and enterococcal populations for potential virulence factors. Microbiol Res. 2008:159(9-10):595-601. doi: 10.1016/j.resmic.2008.09.001.
- Gámez-Valdez JS, García-Mazcorro JF, Montoya-Rincón AH, Rodríguez-Reyes DL, Jiménez-Blanco G, Rodríguez MA, *et al.* Differential analysis of the bacterial community in colostrum samples from women with gestational diabetes mellitus and obesity. Scientific Reports. 2021;11(1): 24373. doi: 10.1038/s41598-021-03779-7.
- Xie W, Zhang H, Ni Y, Peng Y. Contrasting diversity and composition of human colostrum microbiota in a maternal cohort with different ethnic origins but shared physical geography (Island scale). Front Microbiol. 2022;13:934232. doi: 10.3389/fmicb.2022.934232.
- Chande AT, Nagar SD, Rishishwar L, Mariño-Ramírez L, Medina-Rivas MA, Valderrama-Aguirre AE, *et al.* The impact of ethnicity and genetic ancestry on disease prevalence and risk in Colombia. Front Genet. 2021;12:690366. doi: 10.3389/fgene.2021.690366.
- Nascimento RC, Hochman VG, da Silva CB, do Valle BV, do Amaral Y, Dolinsky M, *et al.* Immediate effect of food intake by the nursing mother on the macronutrient content of colostrum. J Pediatr. 2025. doi: 10.1016/j.jped.2025.03.004.
- Londoño-Sierra DC, Mesa V, Guzmán NC, Bolívar Parra L, Montoya-Campuzano OI, Restrepo-Mesa SL. Maternal Diet May Modulate Breast Milk Microbiota—A Case Study in a Group of Colombian Women. Microorganisms. 2023;11(7):1812. doi: 10.3390/microorganisms11071812
- Damaceno QS, Souza JP, Nicoli JR, Paula RL, Assis GB, Figueiredo HC, et al. Evaluation of potential probiotics isolated from human milk and colostrum. Probiotics Antimicrob Proteins. 2017;9: 371-379. doi: 10.1007/s12602-017-9270-1.
- Tawi F. Advantages and limitations of potential methods for the analysis of bacteria in milk: a review. J Food Sci Technol 53, 42–49 (2016). doi: 10.1007/s13197-015-1993-y.